arXiv:math/0602171v3 [math.OC] 20 Apr 2006

Preference fusion when the number of alternatives exceeds two: indirect scoring procedures[⋆]

P. Yu. Chebotarev[1] and Elena Shamis

Institute of Control Sciences of the Russian Academy of Sciences
65 Profsoyuznaya Str., Moscow 117997, Russia

Abstract

We consider the problem of aggregation of incomplete preferences represented by arbitrary binary relations or incomplete paired comparison matrices. For a number of indirect scoring procedures we examine whether or not they satisfy the axiom of *self-consistent monotonicity*. The class of *win-loss combining scoring procedures* is introduced which contains a majority of known scoring procedures. Two main results are established. According to the first one, every win-loss combining scoring procedure breaks self-consistent monotonicity. The second result provides a sufficient condition of satisfying self-consistent monotonicity.

---

1 Introduction

A good method of preference fusion when there are only two alternatives is the simple majority of judge votes. This has been shown by Condorcet in his famous jury theorem [1]; for a review of further developments see [2,3].

The situation becomes much more complicated when the number of alternatives exceeds two. In this case, the majority of judge preferences may turn out to be intransitive and contain preference cycles. Some statistical approaches

---

[⋆] This work was supported by Russian Foundation for Basic Research Grant No. 96–01–01010. Partial research support from the European Community under Grant ACE–91–R02 is also gratefully acknowledged. We are thankful to J. W. Moon, D. S. Schmerling and J. L. Thompson for their help in our search of literature on indirect scoring procedures.

[1] `pchv@rambler.ru`; Participation in the Workshop on Foundations of Information / Decision Fusion was supported by NSF.

to this problem are developed in the theory of paired comparisons (see [4]) and ranking data (see [5]).

In this paper, we use a normative approach typical of the social choice theory and consider indirect scoring procedures as the methods of preference fusion. These ingenious procedures are mainly developed in such disciplines as applied statistics, scoring of sport tournaments, graph theory, management science, etc. Indirect score of an alternative reflects not only the outcomes of its comparisons with other alternatives, but also the comparison outcomes of those alternatives to which this one has been compared, and this way it may depend on all the available preference data. Still there are very few papers on indirect scoring procedures in the social choice theory. Probably, the reason is that these procedures are not arithmetically ascetic enough, and thereby, according to a widespread opinion, do not correspond to the non-quantitative nature of the social choice problem. Another problem is that they are not easy to describe to the individuals involved in the choice process. However, complexity may be considered to be an advantage in the context of the strategic behavior: complicated procedures are more difficult to manipulate.

Nevertheless, there is a case where the indirect scoring procedures are really needed, namely, the case of incomplete preferences. In this paper, we attempt to demonstrate this case, give a brief critical review of indirect scoring procedures, and compare their properties.

Let us consider the situation where an individual is given a set of alternatives to be compared, but he/she is not an expert in all of them. Suppose she is allowed to compare only those alternatives she is familiar with; moreover, to report only those comparison outcomes she is certain of. In fact, we consider even more general types of individual opinions, namely, each of them is an arbitrary binary relation. To prepare these data for calculating indirect scores, we represent them by *incomplete matrices of paired comparisons* borrowed from the statistical theory of paired comparisons.

The main aim of this paper is to examine whether or not the indirect scoring procedures satisfy the axiom of *Self-Consistent Monotonicity* (*SCM*). It has been introduced in [6] where we used it to explore a series of preference aggregation procedures based on the resolution of discrete optimization problems.

Up to now there has not been a comprehensive review of indirect scoring procedures. However, a lot of information can be found in monographs [4,7–10] and papers [11–16]. For relations to the social choice framework see [17]. An adjacent subject is the analysis of tournament social choice rules (see, e.g., [18,19]).

The paper is organized as follows. After the following section which provides main notation, in Section 3 we give an example demonstrating the specific character of aggregating incomplete preferences. Section 4 contains the statement and justification of SCM. Section 5 demonstrates that if a procedure is *based on individual scores*, it break SCM. Sections 6, 8 and 10 give a review of indirect scoring procedures known from the literature. Section 7 introduces the class of *win-loss combining scoring procedures* and establishes that these procedures fail to satisfy SCM. Section 9 provides a sufficient condition for SCM, and Section 11 discusses two other axioms.

2 Notation

Suppose $J = \{1, \dots, n\}$ is a set of alternatives to be compared. There are m individuals (judges, voters, etc.), and each of them reports his/her preferences. Concerning any pair (i, j) of alternatives, an individual may give one of the following four responses: (a) "i is better than j", (b) "j is better than i", (c) "i and j are equivalent", and (d) "I do not report my opinion on this pair." Using the responses of the pth individual, an *incomplete paired comparison matrix* $A^{(p)} = [a_{ij}^{p}]$ is filled out. Its entries a_{ij}^{p} and a_{ji}^{p} correspond to the comparison outcome of i and j as follows:

if the pth individual said that

– i is better than j, then $a_{ij}^{p} = 1$, $a_{ji}^{p} = 0$;
– i and j are equivalent, then $a_{ij}^{p} = a_{ji}^{p} = 1/2$;

If the individual did not report his/her opinion, a_{ij}^{p} and a_{ji}^{p} remain undefined.

By definition, we set all diagonal entries to be zero: $a_{ii}^{p} = 0$, $i = 1, \dots, n$, $p = 1, \dots, m$. Some authors put $a_{ii}^{p} = 1/2$ or leave a_{ii}^{p} undefined, but this does not change the results significantly.

This way we have an array of incomplete paired comparison matrices $\mathcal{A} = (A^{(1)}, \dots, A^{(m)})$ which is referred to as a *profile of individual preferences.* A *scoring* (*rating*) *procedure* is a function $\mathcal{S}$ from the set of all profiles (or its subset) into $\mathbb{R}^n$, where s_i, the ith component of the resulting column vector $\mathbf{s} = (s_1, \dots, s_n)^\top$ is interpreted as a *score of the alternative* i. All the scoring procedures considered in this paper are assumed to be *neutral* (any reindexing of the alternatives preserves their scores) and *anonymous* (any reindexing of the individuals preserves the scores of the alternatives). The term "aggregation of incomplete preferences" will mean here nothing but rating the alternatives by the scores $s_1, \dots, s_n$: the greater is a score, the more socially preferred is the alternative.

3 An Example

The problem of aggregation of incomplete preferences has a specific flavor which cannot be revealed in dealing with complete preferences. To capture it consider the following example.



Fig. 1. a, b, c: Preferences of nine individuals; d: Combined preferences.

Suppose there are four alternatives, $J = \{1, 2, 3, 4\}$, and nine individuals (judges). Three judges are familiar with alternatives 1 and 2 and they all feel that 1 is better (Fig. 1.a where an arrow from 1 to 2 shows that 1 is preferred to 2). Three judges think that 1 and 3 are equivalent (Fig. 1.b where line segments without arrows designate equivalencies). Finally, three judges are familiar with 2 and 4, and they all consider these alternatives of equal quality (Fig. 1.c). The preferences of these nine individuals are incorporated in Fig. 1.d. Can we say anything about the comparative quality of 3 and 4? Although no judge is familiar with 3 and 4 simultaneously, it is plausible that 3 is better than 4. Note that 3 and 4 have the same comparison outcomes: three equivalencies; they differ only in the alternatives to which they were compared. The "opponent" of 3 is probably better than that of 4 and due to this 3 may be estimated higher than 4.

Thus, in aggregating incomplete preferences we should take into account "quality (strength) of the opponents" or "caliber of the opposition" or "schedule difficulty". These terms came from sport, and we see that there is a similarity between the problem of aggregation of incomplete preferences and the problem of rating the participants of an incomplete tournament (note that the extent of this similarity is a distinct and interesting question). Moreover, in the following section this sport analogy will be exploited to explain the meaning of Self-Consistent Monotonicity, the axiom which is discussed throughout the paper. Sometimes the comparison outcome where i is preferred to j will be called a "*win*" of i and a "*loss*" of j.

4 Self-Consistent Monotonicity

First, let us consider one more example. Suppose we have the incomplete sport tournament whose fragment containing all game results of i and j is shown in Fig. 2. Suppose we know that a is stronger than b, c is stronger than d, and e is stronger than f. Then the results of i are definitely better than those of j. Indeed, a loss to a is more pardonable than that to b, a win over c is more honorable than that over d, and a win over e is more valuable than a draw with f. Besides, i has three extra wins and j has two extra losses, which intensify the advantage of i over j.



Fig. 2. An illustration to self-consistent monotonicity.

In fact, the following self-consistent monotonicity axiom requires that in such situations the score of i should be greater than the score of j. The only point is the meaning of preconceptions like "we know that a is stronger than b". Self-consistent monotonicity applies to scoring procedures, and here "we know that a is stronger than b" signifies "*this* scoring procedure gives a a greater score than b." This makes clear why this kind of monotonicity is called "self-consistent".

In the following statement of self-consistent monotonicity, we use the term "multiset". Its difference from set is that multiset may contain the same element in several copies. This concept is needed here since the outcomes of comparisons with different alternatives may coincide. The elements of multisets will be written within angle brackets.

Self-consistent monotonicity (SCM). *A scoring procedure $\mathcal{S}$ is* Self-Consistently Monotonic *if for any set of alternatives J, any profile $\mathcal{A}$, and any alternatives $i, j \in J$ it satisfies the following condition.*

Suppose $U_i = \langle a_{ik}^p \mid k, p \rangle$ and $U_j = \langle a_{j\ell}^q \mid \ell, q \rangle$ are the multisets of the compari-

son outcomes of alternatives i *and* j*, respectively. Suppose that* U_i *can be split into* U_i^I *and* U_i^{II}*, and* U_j *can be split into* U_j^I *and* U_j^{II} *in such a way that*

(*i*) $a_{ik}^p \in U_i^I$ *implies* $a_{ik}^p = 1$, $a_{j\ell}^q \in U_j^I$ *implies* $a_{j\ell}^q = 0$;
(*ii*) *there exists a one-to-one correspondence* π *of* U_i^{II} *onto* U_j^{II} *such that* $\pi(a_{ik}^p) = a_{j\ell}^q$ *implies* $a_{ik}^p \geq a_{j\ell}^q$ *and* $s_k \geq s_\ell$.

Then $s_i \geq s_j$.

Moreover, if in addition U_i^I *is nonempty or* U_j^I *is nonempty or at least one inequality in (ii) is strict at least once, then* $s_i > s_j$.

The application of SCM to a pair of alternatives (i, j) will be called *confrontation of* i *and* j.

It turns out that many well-known indirect scoring procedures fail to satisfy self-consistent monotonicity. In the following two sections we try to explore the features of these procedures that cause them not to satisfy SCM. The results are presented in Proposition 1 (Section 5) and Theorem 8 (Section 7). Section 9 provides a sufficient condition of SCM (Theorem 12).

5 Procedures Based on Individual Scores

A scoring procedure $\mathcal{S}$ is *based on individual scores* if there exist functions f and δ such that for any profile $\mathcal{A} = (A^{(1)}, \ldots, A^{(m)})$, the corresponding score vector $\mathbf{s}$ can be expressed as $\mathbf{s} = \delta(\mathbf{s}^{(1)}, \ldots, \mathbf{s}^{(m)})$, where $\mathbf{s}^{(p)}$ is a partial score vector depending solely on the comparison matrix of individual p: $\mathbf{s}^{(p)} = f(A^{(p)})$, $p = 1, \ldots, m$.

The most important instance is

$$\mathbf{s} = \sum_{p=1}^{m} \mathbf{s}^{(p)},$$

forming (in case of complete preferences) a class of procedures which satisfy the recent axiomatics by Myerson [20].

Proposition 1 *There are scoring procedures based on individual scores that satisfy SCM on complete preferences but no such procedure satisfies SCM for incomplete preferences.*

PROOF. To prove the first statement, it suffices to consider the row sum procedure, which in case of complete preferences has the form

$$
\begin{aligned}
s_i^{(p)} &= \sum_{j=1}^{n} a_{ij}^{p}, \qquad i = 1, \ldots, n, \quad p = 1, \ldots, m, \\
\mathbf{s} &= \sum_{p=1}^{m} \mathbf{s}^{(p)} .
\end{aligned}
$$

This procedure obviously satisfies SCM.

The second statement can be proved by considering our first example of Fig. 1. Let $p = 1, 2, 3$ be the numbers of individuals in Fig. 1.a, $p = 4, 5, 6$ in Fig. 1.b, and $p = 7, 8, 9$ in Fig. 1.c. By neutrality and anonymity of $\mathcal{S}$,

$$
\begin{aligned}
s_3^{(1)} &= s_3^{(2)} = s_3^{(3)} = s_4^{(1)} = s_4^{(2)} = s_4^{(3)}, \\
s_3^{(4)} &= s_3^{(5)} = s_3^{(6)} = s_4^{(7)} = s_4^{(8)} = s_4^{(9)}, \\
s_4^{(4)} &= s_4^{(5)} = s_4^{(6)} = s_3^{(7)} = s_3^{(8)} = s_3^{(9)} .
\end{aligned}
$$

Hence, $s_3 = \sum\limits_{p=1}^{9} s_3^{(p)} = \sum\limits_{p=1}^{9} s_4^{(p)} = s_4$.

Confronting 3 and 4 by SCM (see Fig. 1.d), we deduce $s_1 = s_2$. But now, confronting 1 and 2, we get contradiction. $\Box$

In words, these procedures generally break SCM because the result of confrontation between two alternatives may be only determined by the preferences of the *whole* board of judges.

6 Aggregate Scoring Procedures

Hereafter we will consider scoring procedures that cannot be reduced to individual scores. Such indivisible procedures can be called *aggregate*. A large variety of them are representable through the matrix $A = [a_{ij}]$ of total comparison outcomes on the pairs of alternatives:

$$
a_{ij} = \begin{cases} 0, & \text{if } a_{ij}^{p} \text{ is undefined for every } p, \\ \sum\limits_{p} a_{ij}^{p}, & \text{otherwise.} \end{cases} \tag{1}
$$

Consider the scores of the form:

$$
s_i = \sum_{j=1}^{n} f(a_{ij}, x_j), \quad i = 1, \ldots, n, \tag{2}
$$

where x_j $(j = 1, \dots, n)$ is some estimate of the alternative j, f is a function nondecreasing in both a_{ij} and x_j. Sometimes averaged scores of the form

$$s_i = \frac{1}{m_i} \sum_{j=1}^{n} f(a_{ij}, x_j), \quad i = 1, \dots, n \tag{3}$$

are proposed, where m_i is the total number of comparisons of i (see, e.g., [16]). For simplicity, we do not consider this modification here, however, it is covered by Theorem 8 below.

6.1 Wei's Procedure

The most popular form of f in Eqs (2) is the product:

$$s_i = \sum_{j=1}^{n} a_{ij}\, x_j, \quad i = 1, \dots, n. \tag{4}$$

Regarding the choice of the form of x_j, perhaps the most attractive idea is to relate x_j and s_j directly. However, if we set $x_j = s_j$, $j = 1, \dots, n$, the homogeneous system of linear equations (4) may have only a trivial solution $s_1 = \cdots = s_n = 0$. A minor modification is to define x_j to be proportional to s_j:

$$x_j = s_j / \lambda, \quad j = 1, \dots, n,$$

not specifying λ *a priori*. Then

$$\lambda s_i = \sum_{j=1}^{n} a_{ij}\, s_j, \quad i = 1, \dots, n, \tag{5}$$

or in the matrix form,

$$\lambda \mathbf{s} = A\, \mathbf{s}, \tag{6}$$

thus λ is an eigenvalue and $\mathbf{s}$ an eigenvector of A.

This scoring procedure was proposed by Wei [21] and became well-known after Kendall's [22] paper. Matrix A generally has several eigenvalues, each having its own subspace of eigenvectors. Only one solution is taken, and this solution possesses special properties. Let us suppose that the preferences are *indivisible*, i.e., the set of alternatives J cannot be split into two nonempty parts J_1 and J_2 such that for no alternatives $j \in J_2$ and $i \in J_1$, $a_{ji} > 0$. Then, by the Perron–Frobenius theorem, the largest in absolute value eigenvalue of

A is positive, the corresponding subspace of eigenvectors is one-dimensional, and these eigenvectors have the same sign of all terms. Just the normalized positive eigenvector $\mathbf{s}$ of this type is taken as a vector of scores in the Wei procedure (which applies to indivisible preferences).

Wei's eigenvector can be obtained iteratively. Consider the following sequence of score vectors:

$$\begin{aligned}
&\mathbf{s}^1 = A\,\mathbf{1}, \quad \text{where} \;\; \mathbf{1} = (1,\ldots,1)^\top;\\
&\mathbf{s}^2 = A\,\mathbf{s}^1 = A^2\,\mathbf{1};\\
&\qquad \cdots\cdots\\
&\mathbf{s}^k = A\,\mathbf{s}^{k-1} = A^k\,\mathbf{1};\\
&\qquad \cdots\cdots
\end{aligned}$$

It follows from the Perron–Frobenius theorem that in case of indivisible profiles, the normalized sequence $(\mathbf{s}^k)$ converges to Wei's score vector:

$$\frac{\mathbf{s}^k}{\sum\limits_{i=1}^{n} s_i^k} = \frac{1}{\sum\limits_{i=1}^{n} s_i^k} A^k\,\mathbf{1} \xrightarrow[k\to\infty]{} \mathbf{s}\,. \tag{7}$$

It is a simple fact that s_i^k is the number of k-length paths from i to all vertices in the preference multidigraph. Thus, in Wei's procedure, alternatives are compared by the number of very long ("infinitely-long") paths diverging from them in the preference multidigraph. In Subsection 6.3 we shall consider a scoring procedure where longer paths are accounted with smaller weights.

Proposition 2 *If indivisible preferences are complete, Wei's scoring procedure satisfies SCM, but it breaks SCM for incomplete indivisible preferences.*

PROOF. To prove the first statement, confront arbitrary i and j by SCM:

$$s_i = \frac{1}{\lambda} \sum_{k=1}^{n} a_{ik}\, s_k, \tag{8}$$

$$s_j = \frac{1}{\lambda} \sum_{k=1}^{n} a_{jk}\, s_k, \tag{9}$$

and assume that some partitions of U_i and U_j described in self-consistent monotonicity exist. Then U_i^{II} induces on the right-hand side of Eq. (8) a no lesser (respectively, greater in the strict case) sum of terms than U_j^{II} induces on the right-hand side of Eq. (9). The same can be said of U_i^I and U_j^I. Therefore, $s_i \geq s_j$ ($s_i > s_j$ in the strict case).

To prove the second statement, consider the cumulative preference digraph depicted in Fig. 3.



Fig. 3. Preferences in the proof of Proposition 2.

The comparison outcomes of alternative 5 are equivalencies with alternatives 1, 2, 3 and 4. Alternative 5 is introduced only to make the preferences indivisible. Consider the equations with s_2 and s_3 on the left-hand sides:

$$\begin{aligned}\lambda s_2 &= s_4 + s_5/2,\\ \lambda s_3 &= s_4 + s_5/2.\end{aligned}$$

Hence, $s_2 = s_3$, which contradicts SCM, since 3 has an extra "loss" ($a_{31} = 0,\ a_{13} = 1$). □

The example used in the proof of Proposition 2 reveals one important feature of Wei's procedure as applied to incomplete preferences. Namely, this procedure bases itself only on "wins" and does not take into account "losses". More precisely, it does not distinguish "losses" from missing comparisons (and precisely this contradicts SCM). A possible way to make the procedure more balanced is to also consider a dual procedure which is based on "losses", and then to combine the resulting scores. This has been done by Hasse [23] and Ramanujacharyulu [24].

6.2 The Procedures by Hasse and Ramanujacharyulu

Let $\mathbf{w}$ and $\mathbf{l}$ denote the right and left eigenvectors of matrix A, corresponding to its maximal eigenvalue λ :

$$\lambda w_i = \sum_{j=1}^{n} a_{ij} w_j, \quad i = 1, \ldots, n, \tag{10}$$

$$\lambda \ell_i = \sum_{j=1}^{n} a_{ji}\, \ell_j, \quad i = 1, \ldots, n, \tag{11}$$

Here $\mathbf{w} = (w_1, \ldots, w_n)$ is the vector of Wei's *win-scores*, which is not sensitive to the distinction between "losses" and missing comparisons; $\mathbf{l} = (\ell_1, \ldots, \ell_n)$ is the vector of *loss-scores*, which does not distinguish "wins" and missing comparisons. The matrix form of Eqs (11) is

$$\lambda\, \mathbf{l} = B\, \mathbf{l}, \tag{12}$$

where $B = A^\top$ is the transpose of A.

Hasse [23] proposed to combine $\mathbf{w}$ and $\mathbf{l}$ into the ultimate scores:

$$s_i = w_i - \ell_i, \quad i = 1, \ldots, n. \tag{13}$$

Another procedure proposed by Ramanujacharyulu [24] has scores given by:

$$s_i = w_i / \ell_i, \quad i = 1, \ldots, n. \tag{14}$$

These two procedures are close but may produce different rankings, since the first one registers the absolute differences between win-scores and loss-scores, whereas the second is based on the relative differences. Both procedures are applicable to indivisible preferences and turn out to break self-consistent monotonicity. We do not formulate the corresponding statement, because there are many possible modifications of such procedures, and a more reasonable thing to do is to isolate their common features and then to prove a more general theorem. To this end, in the following two subsections we consider two other families of indirect scoring procedures, and then proceed with a theorem.

6.3 The Procedure of Katz–Thompson–Taylor

As we have seen, Wei's scores rank order the alternatives according to the number of very long paths diverging from them in the preference multidigraph. Katz [25] and then Thompson [26] and Taylor [27] proposed a scoring procedure in which the long paths play just a correcting role, whereas the short paths are taken into account with greater weights. Namely, Thompson introduced the scores of the form

$$\mathbf{w} = (A + \varepsilon A^2 + \varepsilon^2 A^3 + \cdots)\, \mathbf{1}, \tag{15}$$

where ε is a small positive parameter. If $\varepsilon < r^{-1}$ where r is the spectral radius of A, then the series (15), which in fact is a geometric progression, converges and

$$\mathbf{w} = A(I - \varepsilon\, A)^{-1}\, \mathbf{1}, \tag{16}$$

where I is the identity matrix.

Thompson shows that his normalized vector $\mathbf{w}$ tends to Wei's eigenvector as ε approaches r^{-1} from below. Besides, he gives a game-theoretical interpretation of these scores comparable with the approach by Laffond, Laslier and Lebreton [28].

The paper by Taylor is one of the few social choice articles devoted to indirect scoring procedures. He investigated sequential voting by the committee and considered Hamiltonian sequences of winning alternatives, which led him to a scoring procedure very close to (15). In fact, Taylor used another matrix $C = [c_{ij}]$ instead A, where

$$c_{ij} = \begin{cases} a_{ij} - a_{ji}, & \text{if } a_{ij} - a_{ji} \geq 0, \\ 0, & \text{otherwise.} \end{cases}$$

Since for the preference structure we use in the proof of Theorem 8 in Section 7, A and C coincide, the statement of this theorem is valid for Taylor's procedure.

As above, $\mathbf{w}$ in Eqs (15) and (16) is a vector of *win-scores*, and it is worth being supplemented by the corresponding *loss-scores*. So we may consider a pair of vectors $(\mathbf{w}, \mathbf{l})$ where

$$\mathbf{l} = (B + \varepsilon\, B^2 + \varepsilon^2\, B^3 + \cdots)\, \mathbf{1}, \tag{17}$$

and combine $\mathbf{w}$ and $\mathbf{l}$ as in the procedures of Hasse and Ramanujacharyulu:

$$s_i = w_i - \ell_i, \quad i = 1, \ldots, n$$

or

$$s_i = w_i / \ell_i, \quad i = 1, \ldots, n.$$

The question we are interested in is whether or not these procedures satisfy self-consistent monotonicity. The answer is negative. After considering one more family of scoring procedures, we give a theorem that will clarify this point to some extent.

6.4 Directed Tree Procedure by Daniels–Ushakov–Goddard–Levchenkov

Let c_i^+ and c_i^- be the total "win" and total "loss" of i, respectively:

$$\begin{aligned} c_i^+ &= \sum_{j=1}^{n} a_{ij}, \\ c_i^- &= \sum_{j=1}^{n} a_{ji}, \quad i = 1, \ldots, n. \end{aligned} \tag{18}$$

Daniels [11] and other writers [29–33] proposed, with various motivations, the scores $w_1, \ldots, w_n$ that satisfy

$$w_i = \frac{1}{c_i^-} \sum_{j=1}^{n} a_{ij} w_j, \quad i = 1, \ldots, n. \tag{19}$$

The only difference of this system of equation from that of Wei is that λ is replaced by c_i^-. This is possible because the matrix $[a_{ij}/c_i^-]$ always has a unit eigenvalue. Scores w_i have several interesting interpretations listed below.

(i) w_i is proportional to the number of directed trees diverging from i in the preference multidigraph [34];
(ii) w_i are proportional to the final probabilities of the "winning Markov chain" [29,30,35,32,33,36];
(iii) $\mathbf{w}$ is the vector of "fair bets" in the model of Moon and Pullman [12].

Also, in [37–39] this procedure is analyzed in the social choice framework, and in [40,41] its strategic properties are touched upon.

Exactly as before, we may consider the corresponding *loss-scores* mentioned by Daley [35]:

$$\ell_i = \frac{1}{c_i^+} \sum_{j=1}^{n} a_{ji} \ell_j, \quad i = 1, \ldots, n \tag{20}$$

and define the ultimate score vector as a combination of $\mathbf{w}$ and $\mathbf{l}$, e.g.,

$$s_i = w_i - \ell_i, \quad i = 1, \ldots, n$$

or

$$s_i = w_i / \ell_i, \quad i = 1, \ldots, n.$$

The procedures of this family break SCM, and now we are in a position to specify a more general class of indirect scoring procedures that do not obey SCM. All the above procedures are included.

7 A General Class of Procedures That Fail to Satisfy Self-Consistent Monotonicity

In this section we define *win-loss indices* and *win-loss combining procedures* and prove that such procedures break self-consistent monotonicity.

In the following Definitions 3–5, profile $\mathcal{A}$ is fixed; "vector" means n-component column vector.

Definition 3 (*recursive*). Win-loss index of finite order.

1°. *A pair of vectors* $(\mathbf{w}, \mathbf{l})$ *is a* win-loss index of zero order *if* $\mathbf{w} = \mathbf{l} = (1, \ldots, 1)^\top$.

2°. *A pair of vectors* $(\mathbf{w}, \mathbf{l})$ *is a* win-loss index of order k *if there exists a win-loss index* $(\mathbf{w}', \mathbf{l}')$ *of order* $k-1$ *and functions* f, g, α *and* β *such that*

$$\begin{aligned} w_i &= \frac{1}{\alpha(c_i^+, c_i^-)} f(W_i'), \\ \ell_i &= \frac{1}{\beta(c_i^-, c_i^+)} g(L_i'), \quad i = 1, \ldots, n, \end{aligned} \tag{21}$$

where $W_i' = \langle (a_{i1}, w'_1), (a_{i2}, w'_2), \ldots, (a_{in}, w'_n) \rangle$, $L_i' = \langle (a_{1i}, \ell'_1), (a_{2i}, \ell'_2), \ldots, (a_{ni}, \ell'_n) \rangle$, $i = 1, \ldots, n$; *angle brackets designate multisets.*

W_i' *and* L_i' *are respectively called the* win-performance *and the* loss-performance *of* i *corresponding to* $(\mathbf{w}', \mathbf{l}')$.

One can see that in the Directed tree procedure of Subsection 6.4, $\alpha(c_i^+, c_i^-)$ is c_i^-; $\beta(c_i^-, c_i^+)$ is c_i^+; f and g are sums of products. An essential difference is that in the Directed tree procedure, w_i and ℓ_i are related with the win-performance and loss-performance corresponding to the same pair of vectors $(\mathbf{w}, \mathbf{l})$, not to a win-loss index of the previous order. Thus, Eqs (19) and (20) do not fit Definition 3. That pair of vectors $(\mathbf{w}, \mathbf{l})$ satisfies the following definition.

Definition 4 Win-loss index of infinite order. *A pair of vectors* $(\mathbf{w}, \mathbf{l}) = ((w_1, \ldots, w_n)^\top, (\ell_1, \ldots, \ell_n)^\top)$ *is a* win-loss index of infinite order *if there exist functions* f, g, α *and* β *such that*

$$\begin{aligned} w_i &= \frac{1}{\alpha(c_i^+, c_i^-)} f(W_i), \\ \ell_i &= \frac{1}{\beta(c_i^-, c_i^+)} g(L_i), \quad i = 1, \ldots, n, \end{aligned} \tag{22}$$

where $W_i = \langle (a_{i1}, w_1), (a_{i2}, w_2), \ldots, (a_{in}, w_n) \rangle$ *and* $L_i = \langle (a_{1i}, \ell_1), (a_{2i}, \ell_2), \ldots, (a_{ni}, \ell_n) \rangle$, $i = 1, \ldots, n$.

The following definition introduces win-loss indices without order.

Definition 5 Win-loss index. *A pair of vectors* $(\mathbf{w},\mathbf{l}) = ((w_1,\ldots,w_n)^\top, (\ell_1,\ldots,\ell_n)^\top)$ *is a* win-loss index *if there exists a sequence* $((w^1,\ell^1),(w^2,\ell^2),\ldots)$ *of win-loss indices of finite or infinite order and functions* φ *and* ψ *such that for* $i=1,\ldots,n$*,*

$$\begin{aligned} w_i &= \varphi(w_i^1,\ell_i^1,w_i^2,\ell_i^2,\ldots),\\ \ell_i &= \psi(w_i^1,\ell_i^1,w_i^2,\ell_i^2,\ldots). \end{aligned} \tag{23}$$

The components of $\mathbf{w}$ and $\mathbf{l}$ are referred to as *win-scores* and *loss-scores*, respectively (note that they depend on both w-components and ℓ-components of the sequence elements). The next (and last) definition introduces a scoring procedure that combines win-scores and loss-scores of a win-loss index.

Definition 6 Win-loss combining scoring procedure. *Scoring procedure* $\mathcal{S}$ *is a* win-loss combining scoring procedure *if for any profile* $\mathcal{A}$ *there exists a win-loss index* $(\mathbf{w},\mathbf{l})$ *and a function* h *such that*

$$s_i = h(w_i,\ell_i), \quad i=1,\ldots,n.$$

Proposition 7 *All scoring procedures that generate score vectors of Section 6 or any their iterative approximations constructed as in* 2° *of Definition 3, are win-loss combining scoring procedures.*

The proof of Proposition 7 is straightforward. Among other win-loss combining scoring procedures, let us mention the elaborate procedure by David (see [42,4,15], and procedures by Cowden [43] and Ginovker [44]. The latter two authors also proposed other procedures which will be mentioned below. An interesting discussion of related topics can be found in [16].

Theorem 8 *Every win-loss combining scoring procedure defined on the set of indivisible preference profiles breaks self-consistent monotonicity.*

PROOF. Assume that there exists a win-loss combining scoring procedure that satisfies SCM on the set of indivisible preference profiles. Suppose that $(\mathbf{w},\mathbf{l})$ is its win-loss index for the profile with the preference digraph shown in Fig. 4 and that $s_i = h(w_i,\ell_i)$, $i=1,\ldots,n$.

Lemma 9 *In the above assumptions,* (i) $\ell_2=\ell_3$*;* (ii) $w_2=w_3$.

Proof of Lemma 9. First, assume that $(\mathbf{w},\mathbf{l})$ is a win-loss index of kth order. If $k=0$, (i) and (ii) of Lemma 9 hold. Note also that $w_6=w_7$. If $k>0$, there



Fig. 4. Preferences in the proof of Theorem 8.

exists a win-loss index $(\mathbf{w}', \mathbf{l}')$ of $(k-1)$th order and functions f, g, α and β such that Eqs (21) hold for $i = 1, \ldots, 7$. In particular,

$$w_6 = \tfrac{1}{\alpha(1,2)} f\left(\langle (1, w'_1), (0, w'_2), \ldots, (0, w'_7) \rangle\right) = w_7.$$

(i) Similarly, $\ell_2 = \ell_3$.
(ii) $w_2 = \tfrac{1}{\alpha(1,1)} f\left(\langle (0, w'_1), \ldots, (0, w'_5), (1, w'_6), (0, w'_7) \rangle\right)$,
$w_3 = \tfrac{1}{\alpha(1,1)} f\left(\langle (0, w'_1), \ldots, (0, w'_5), (1, w'_7), (0, w'_6) \rangle\right)$.

In the latter formula we interchanged the two last elements of the multiset W'_3. As we have seen above, for every win-loss index of finite order, $w_6 = w_7$, hence $w'_6 = w'_7$. Therefore, $w_2 = w_3$.

For win-loss indices of infinite order, (i) and (ii) are proved similarly.

Suppose now that $(\mathbf{w}, \mathbf{l})$ is an arbitrary win-loss index. Then there exists a sequence of win-loss indices of finite or infinite order $((w^1, \ell^1), (w^2, \ell^2), \ldots)$ and functions φ and ψ such that

$$\begin{aligned} w_i &= \varphi(w_i^1, \ell_i^1, w_i^2, \ell_i^2, \ldots), \\ \ell_i &= \psi(w_i^1, \ell_i^1, w_i^2, \ell_i^2, \ldots), \qquad i = 1, \ldots, 7. \end{aligned}$$

Since for every $t = 1, 2, \ldots,$ $\ell_2^t = \ell_3^t$ and $w_2^t = w_3^t$, we get $\ell_2 = \ell_3$ and $w_2 = w_3$. Lemma 9 is proved.

Since $s_2 = h(w_2, \ell_2)$ and $s_3 = h(w_3, \ell_3)$, Lemma 9 implies $s_2 = s_3$.

Let us contrast alternatives 2 and 3 by self-consistent monotonicity. It follows that if either $s_6 > s_7$ or $s_7 > s_6$, then $s_2 = s_3$ is impossible. Hence, $s_6 = s_7$. Now, contrasting alternatives 4 and 5, we get $s_4 > s_5$.

Finally, contrasting 6 and 7 and using the proved statements $s_2 = s_3$ and $s_4 > s_5$ we get $s_6 > s_7$, which contradicts $s_6 = s_7$ obtained above. Thus, our assumption is wrong, and self-consistent monotonicity is broken. $\square$

Theorem 8 can be explained as follows. All functions in Definitions 3–6 are arbitrary, we impose no constraints on them. The only non-arbitrariness was that “wins” and “losses” were *treated separately* (whereas “draws” influenced both win-scores and loss-scores). This separation was not complete, because the expressions (21) and (22) for win-scores w_i also contained the total losses c_i^- and vice versa; such a cross-dependence extended to win-indices and loss-indices (23) too. In spite of that, as can be shown, a win of alternative j over alternative k can affect the win-score of $i \neq k$ only if there exists a directed path in the preference multidigraph from i to j or from i to k. Similarly, this win of j over k can affect the loss-score of $i \neq j$ only if there exists a directed path from j or from k to i. Therefore the ultimate scores are sensitive only to directed paths of these two kinds. As can be concluded from the proof of Theorem 8, self-consistent monotonicity implies the sensitivity of the ultimate scores to the influences that spread not only through directed paths, but also via paths with altering directions of arrows.

8 Win-Loss Unifying Procedures

In this section we consider two scoring procedures that do not separate wins and losses in the score equations, treating them *uniformly*. Another important difference from the procedures of Section 6 is that here the function combining the comparison outcomes and the estimates of the opponents (function f in Eqs (2)) has an additive (instead of productive) form. It is worth noting that if the score of i is represented through multiplicative terms such as $a_{ij}\, s_j$, then the influence of s_j upon the score of i depends on a_{ij}: the greater a_{ij}, the stronger this influence. In the extreme case where $a_{ij} = 0$, s_j is not taken into account at all. An additive form of scores, on the contrary, causes a uniform influence of s_j upon s_i, regardless of a_{ij}. This appears more reasonable.

8.1 The Least-Squares Procedure

This procedure, first proposed by Smith [45] and Gulliksen [46] and then investigated in [47–52, etc.], constructs a mean square approximation of the comparison outcomes by the differences between the desired scores:

$$\underset{\mathbf{s}}{\text{minimize}} \sum_{i,j,p} (r_{ij}^{p} - (s_i - s_j))^2, \tag{24}$$

where

$$r_{ij}^{p} = a_{ij}^{p} - a_{ji}^{p} \tag{25}$$

is a skew-symmetric modification of the comparison outcomes; r_{ij}^{p} are undefined whenever a_{ij}^{p} and a_{ji}^{p} are undefined; the sum extends over those i, j and p for which r_{ij}^{p} is definite.

Partial differentiation reduces the problem (24) to the system of linear equations

$$s_i = \frac{1}{m_i} \sum_{j,p} (s_j + r_{ij}^{p}), \quad i = 1, \dots, n, \tag{26}$$

where $m_i = c_i^{+} + c_i^{-}$ is the number of comparisons of i.

Note some parallelism of this system of equations and that of (19). A multiplicative counterpart of Eqs (26) is considered in [16] and fits (3).

This procedure is applicable not only to indivisible preference profiles, but to all profiles with connected preference multigraph. Under this condition, the rank of the system (26) is $n-1$, and scores $s_1, \dots, s_n$ can be found up to an additive constant. For the complete preferences, this procedure reduces to that of row sums, and it is rather well-performing for incomplete numerical (weighted) preferences. However, in the case of discrete incomplete preferences which we consider here, it can produce some unnatural results (see the proof of the following proposition).

Proposition 10 *The Least squares procedure breaks self-consistent monotonicity.*

PROOF. Consider the preference multidigraph shown in Fig. 5.

The least-squares score of 2 is greater than that of 1, in spite of that 1 has an extra win. This breaks self-consistent monotonicity. □



Fig. 5. Preferences in the proof of Proposition 10.

This proof can be commented as follows. The Least-squares procedure better fits numerical preferences; it punishes 1 for the win over 4 because it “expects” that if 1 beats 3 and 3 beats 4 with the same “intensity”, 1 should beat 4 with a greater intensity. The following procedure eliminates this flaw.

8.2 The Generalized Row Sum Procedure

This procedure [53,54] can be considered as a Bayesian modification of the previous one. It is derived axiomatically and has Markov chain and graph theoretic interpretations [55,56]. The scores satisfy the system of equations

$$s_i = \varepsilon \sum_{j,p} (\gamma r_{ij}^p - (s_i - s_j)),\ i = 1, \ldots, n, \tag{27}$$

where ε is a positive parameter, $\varepsilon \leq (m(n-2))^{-1}$, and $\gamma = mn + \varepsilon^{-1}$.

Proposition 11 *The Generalized row sum procedure satisfies Self-Consistent Monotonicity.*

The proof is given in [6]. The form of these scores suggests what kind of procedures satisfies self-consistent monotonicity. A class of such procedures is described in the following section.

9 A Sufficient Condition of Self-Consistent Monotonicity

Theorem 12 *Suppose that a scoring procedure $\mathcal{S}$ is such that there exists a function f defined on finite nonempty multisets of real triples and possessing of the following properties.*

(i) for every $\mathcal{A}$,

$$f(\langle(a_{ij}^p, s_i, s_j) \mid j, p\rangle) = 0, \quad i = 1, \ldots, n. \tag{28}$$

(ii) f is
- *increasing in every a_{ij}^p,*
- *increasing in every s_j ($j \neq i$),*
- *decreasing in s_i.*

(iii) Let $V_i = \langle(a_{ij}^p, s_i, s_j) \mid j, p\rangle$. For every $\mathcal{A}$ and for every $i, k \in J$,
if $(1, s_i, s_k) \in V_i$, then $f(V_i \setminus \langle(1, s_i, s_k)\rangle) < f(V_i)$;
if $(0, s_i, s_k) \in V_i$, then $f(V_i \setminus \langle(0, s_i, s_k)\rangle) > f(V_i)$.

Then scoring procedure $\mathcal{S}$ satisfies self-consistent monotonicity.

PROOF. Assume that the relation between i and j described in the statement of self-consistent monotonicity holds, but $s_i < s_j$ (respectively, $s_i \leq s_j$ in the strict case). Then, by (ii) and (iii) of Theorem 12, the left-hand side of the ith equation in the system (28) exceeds the left-hand side of the jth equation, and hence they both cannot be equal to zero. This contradiction proves the theorem. □

In [57] a modification of SCM is proposed which is met by those and only those scoring procedures that have the implicit form (i)–(ii) of Theorem 12.

10 Four Procedures Satisfying SCM

Table 1 lists four procedures that satisfy self-consistent monotonicity.

Table 1
Scoring procedures that satisfy SCM.

Procedure	Domain*	$f(\cdot)$ of Theorem 12
Zermelo [58], Bradley&Terry [59],...	ID	$\sum_{j,p}\left(a_{ij}^p - \frac{s_i}{s_i+s_j}\right)$
Daniels [11], Ginovker [44]	ID	$\sum_{j=1}^{n}\left(a_{ij}\,\frac{s_j}{s_i} - a_{ji}\frac{s_i}{s_j}\right)$
Cowden [43]	ID	$\sum_{j,p}(a_{ij}\, s_j(1-s_i) - a_{ji}\, s_i(1-s_j))$
Generalized row sum procedure [54]	U	$\varepsilon\sum_{j,p}(\gamma r_{ij}^p - (s_i - s_j)) - s_i$

* ID means indivisible preference profiles, U all profiles.

The second column conveys the set of preference profiles to which the corresponding procedure is applicable. Of the four methods, the Generalized row sum procedure is based on the solution of a linear system of equations, three others are reduced to nonlinear systems of equations, which are usually solved with iterative algorithms.

A question arises of how to compare further the procedures that satisfy self-consistent monotonicity. The last section contains two remarks on the possible additional axioms.

11 Concluding Remarks

In a follow-up paper we will continue the axiomatic testing of indirect scoring procedures and turn to their axiomatic derivation. We now give two possible conditions which can supplement self-consistent monotonicity.

11.1 Macrovertex Independence

The main idea of self-consistent monotonicity is sensitivity of the aggregating procedure. However, there exists an independence condition called *Macrovertex independence* that does not contradict SCM and seems rather natural (cf. the discussion after Theorem 8).

A subset M of the set of alternatives J is called a *macrovertex* of the preference multigraph if for every $i, j \in M$ and every $k \in J \setminus M$, $n_{ij} = n_{jk}$, where n_{ik} is the number of comparisons between i and k in $\mathcal{A}$.

Macrovertex independence. *The comparison outcomes of the alternatives within a macrovertex have no influence on the scores of the alternatives outside the macrovertex.*

11.2 Splitting Balance

Let us note that self-consistent monotonicity is not a very restrictive condition, since it leaves room for some preconception. To conclude with what we started, we address our first example again and give it a sport interpretation. Suppose that Fig. 1.d depicts an incomplete chess tournament. Assume that 4 is the world chess champion, 1, 2 and 3 being university students. Our aim is to estimate the strength of the players taking into account prior information.

Most probably, the world champion is the strongest player, and we should allot the highest rank to 4. Player 2 managed to make three draws with the champion. That is great, and we rank 2 second in spite of three his/her losses to 1. Then 1 is the third and 3 the last. Note that this rank order does not contradict SCM! This is what we meant by saying that SCM leaves room for some preconception. There exists a logic (a bit biased) that justifies ranking $(4, 2, 1, 3)$. To avoid this, the following condition may be added.

Splitting Balance. *Suppose the set of alternatives* J *can be split into* J_1 *and* J_2 *such that for no* $i \in J_1$ *and* $j \in J_2$, $a_{ji} > 0$. *Then there exist* $i \in J_1$ *and* $j \in J_2$ *such that* $s_i \geq s_j$.

The objective of this paper was to put together various indirect scoring procedures (mainly, of infinite order) and to subject them to the analysis in the spirit of the social choice theory. We considered the case of incomplete preferences and an axiom we referred to as self-consistent monotonicity. It was established that many of the known indirect scoring procedures are *win-loss combining procedures* and, by Theorem 8 of Section 7, they break self-consistent monotonicity. In our opinion, some of them are applicable in the analysis of social networks, since here self-consistent monotonicity is not so attractive as in preference aggregation (cf. [25,60,61]). For instance, the popularity index of an individual should not strongly depend on his/her own responses.

In Theorem 12 we proposed a sufficient condition for self-consistent monotonicity and then listed four procedures that satisfy it. The analysis of indirect scoring procedures started in this paper is intended to be continued with additional axioms (two possible conditions were mentioned in this section), and with the end goal of axiomatic construction of such procedures.

References

[1] M. J. A. Marquis de Condorcet, “Essai sur l’application de l’analyse à la probabilité des décisions rendues à la pluralité des voix”, Paris, 1785.

[2] B. Grofman and G. Owen, “Review essay: Condorcet models, avenues for further research”, in *Information Pooling and Group Decision Making* (authors, ed.), pp. 93–102, Jai Press, Greenwich, CT, 1986.

[3] S. Berg, “Evaluation of some weighted majority decision rules under dependent voting”, *Mathematical Social Sciences*, Vol. 28, pp. 71–83, 1994.

[4] H. A. David, “The Method of Paired Comparisons”, Griffin, London, Second Edn, 1988.

[5] M. A. Fligner and J. S. Verducci, eds., "Probability Models and Statistical Analyses for Ranking Data", Springer-Verlag, New York, 1993.

[6] P. Yu. Chebotarev and E. Shamis, "Constructing an objective function for aggregating incomplete preferences", in *Econometric Decision Models* (A. Tangian and J. Gruber, eds.), Lecture Notes in Economics and Mathematical Systems, pp. 100–124, Springer-Verlag, Berlin, 1997.

[7] J. W. Moon, "Topics on Tournaments", Holt, Rinehart and Winston, New York, 1968.

[8] A. R. Belkin and M. S. Levin, "Decision Making: Combinatorial Models of Information Approximation", Nauka, Moscow, 1990. (In Russian).

[9] R. Van Blokland-Vogelesang, "Unfolding and Group Consensus Ranking for Individual Preferences", DSWO Press, Leiden, 1991.

[10] W. D. Cook and M. Kress, "Ordinal Information and Preference Structures: Decision Models and Applications", Prentice-Hall, Englewood Cliffs, NJ, 1992.

[11] H. E. Daniels, "Round-robin tournament scores", *Biometrika*, Vol. 56, pp. 295–299, 1969.

[12] J. W. Moon and N. J. Pullman, "On generalized tournament matrices", *SIAM Review*, Vol. 12, pp. 384–399, 1970.

[13] H. A. David, "Ranking the players in a round robin tournament", *Review of International Statistical Institute*, Vol. 39, pp. 137–147, 1971.

[14] D. S. Shmerling, S. A. Dubrovskii, D. Arzhanova, and A. A. Frenkel', "Expert judgments: Methods and applications", in *Statistical Methods of Analysis of Expert Judgments*, Uchen. Zap. Stat., Vol. 29, pp. 290–382, Nauka, Moscow, 1977. (In Russian).

[15] H. A. David and D. M. Andrews, "Nonparametric methods of ranking from paired comparisons", in *Probability Models and Statistical Analyses for Ranking Data* (M. A. Fligner and J. S. Verducci, eds.), pp. 20–36, Springer-Verlag, New York, 1993.

[16] J. P. Keener, "The Perron-Frobenius theorem and the ranking of football teams", *SIAM Review*, Vol. 35, pp. 80–93, 1993.

[17] J. Levin and B. Nalebuff, "An introduction to vote-counting schemes", *Journal of Economic Perspectives*, Vol. 9, pp. 3–26, 1995.

[18] G. Laffond, J.-F. Laslier, and M. Lebreton, "Condorcet choice correspondences: A set-theoretical comparisons", *Mathematical Social Sciences*, Vol. 30, pp. 23–35, 1995.

[19] G. Laffond, J. Lainé, and J.-F. Laslier, "Composition-consistent tournament solutions and social choice functions", *Social Choice and Welfare*, Vol. 13, pp. 75–93, 1996.

[20] R. B. Myerson, "Axiomatic derivation of scoring rules without the ordering assumption", *Social Choice and Welfare*, Vol. 12, pp. 59–74, 1995.

[21] T. Wei, "The Algebraic Foundations of Ranking Theory", Ph.D. Thesis, Cambridge University, 1952.

[22] M. G. Kendall, "Further contributions to the theory of paired comparisons", *Biometrics*, Vol. 11, pp. 43–62, 1955.

[23] M. Hasse, "Über die Behanglung graphentheoretischer Probleme unter Verwendung der Matrizenrechnung", *Wissenschaftliche Zeitschrift Techniscen Universität Dresden*, Vol. 10, pp. 1313–1316, 1961.

[24] C. Ramanujacharyulu, "Analysis of preferential experiments", *Psychometrika*, Vol. 29, pp. 257–261, 1964.

[25] L. Katz, "A new status index derived from sociometric analysis", *Psychometrika*, Vol. 18, pp. 39–43, 1953.

[26] G. L. Thompson, "Lectures on game theory, Markov chains and related topics." Sandia Corporation Monograph SCR–11, 1958.

[27] M. Taylor, "Graph-theoretic approaches to the theory of social choice", *Public Choice*, Vol. 4, pp. 35–47, Spring 1968.

[28] G. Laffond, J.-F. Laslier, and M. Lebreton, "The bipartisan set of a tournament game", *Games and Economic Behavior*, Vol. 15, pp. 182–201, 1993.

[29] I. A. Ushakov, "The problem of choosing the preferable object", *Izv. Akad. Nauk SSSR. Tekhn. Kibernet.*, No. 4, pp. 3–7, 1971. (In Russian).

[30] I. A. Ushakov, "The problem of choosing the preferred element: An application to sport games", in *Management Science in Sports* (R. E. Machol, S. P. Ladany, and D. G. Morrison, eds.), pp. 153–161, North-Holland, Amsterdam, 1976.

[31] S. T. Goddard, "Ranking in tournaments and group decisionmaking", *Management Science*, Vol. 29, pp. 1384–1392, 1983.

[32] V. S. Levchenkov, "Entropy analysis of the majority rule", *Automation and Remote Control*, Vol. 51, pp. 517–524, 1990.

[33] V. S. Levchenkov, "Choice without Arrow's axiom on the independence of irrelevant alternatives", *Doklady Akademii Nauk SSSR*, Vol. 322, pp. 651–655, 1992. (in Russian).

[34] K. A. Berman, "A graph theoretical approach to handicap ranking of tournaments and paired comparisons", *SIAM Journal on Algebraic and Discrete Methods*, Vol. 1, pp. 359–361, 1980.

[35] D. J. Daley, "Markov chains and a pecking order problem", in *Interactive Statistics* (D. McNeil, ed.), pp. 247–254, North-Holland, Amsterdam, 1979.

[36] V. S. Levchenkov, "Choosing by tournament relations", *Doklady Akademii Nauk*, Vol. 336, pp. 324–327, 1994. (In Russian).

[37] V. S. Levchenkov and I. L. Grivko, “Some properties of self-consistent choice”, *Doklady Akademii Nauk*, Vol. 327, pp. 42–47, 1992. (In Russian).

[38] V. S. Levchenkov, “Axiomatic approach to the self-consistent choice”, *Doklady Akademii Nauk*, Vol. 330, pp. 173–176, 1993. (in Russian).

[39] I. L. Grivko and V. S. Levchenkov, “Intrinsic properties of the self-consistent choice rule”, *Automation and Remote Control*, Vol. 55, pp. 689–697, 1994.

[40] V. S. Levchenkov, “Self-consistent description of the manipulation by alternatives”, *Doklady Akademii Nauk*, Vol. 331, pp. 567–570, 1993. (in Russian).

[41] V. S. Levchenkov, “Strategic scheme for group choice”, *Doklady Akademii Nauk*, Vol. 341, pp. 320–323, 1995. (in Russian).

[42] H. A. David, “Ranking from unbalanced paired-comparison data”, *Biometrika*, Vol. 74, pp. 432–436, 1987.

[43] D. J. Cowden, “A method of evaluating contestants”, *American Statistician*, Vol. 29, No. 2, pp. 82–84, 1975.

[44] V. D. Ginovker, “On one approach to evaluating objects from paired comparison results.” VINITI Manuscript No. 3431-81DEP, 1981. (In Russian).

[45] J. H. Smith, “Adjusting baseball standings for strength of teams played”, *American Statistician*, Vol. 10, No. 3, pp. 23–24, 1956.

[46] H. Gulliksen, “A least squares solution for paired comparisons with incomplete data”, *Psychometrika*, Vol. 21, pp. 125–134, 1956.

[47] G. E. Noether, “Remarks about a paired comparison model”, *Psychometrika*, Vol. 25, pp. 357–367, 1960.

[48] P. S. H. Leeflang and B. M. S. Van Praag, “A procedure to estimate relative powers in binary contacts and an application to Dutch football league results”, *Statistica Neerlandica*, Vol. 25, pp. 63–80, 1971.

[49] R. J. Leake, “A method for ranking teams: With an application to college football”, in *Management Science in Sports* (R. E. Machol, S. P. Ladany, and D. G. Morrison, eds.), pp. 153–161, North-Holland, Amsterdam, 1976.

[50] H. F. Kaiser and R. C. Serlin, “Contributing to the method of paired comparisons”, *Applied Psychological Measurement*, Vol. 2, pp. 421–430, 1978.

[51] A. I. Vereskov, “Extended Methods of Regression Analysis in the Problems of Identification of Nonlinear Models”, Ph.D. Thesis, VNIISI, Moscow, 1986. (In Russian).

[52] V. M. Pliner, “On one approach to incomplete paired comparisons”, in *Abstracts of the Fourth All-Union Conference on Applications of Multivariate Statistical Analysis in Economics and Quality Evaluation*, (Tartu), pp. 29–30, 1989. (In Russian).

[53] P. Yu. Chebotarev, “Generalization of the row sum method for incomplete paired comparisons”, *Automation and Remote Control*, Vol. 50, pp. 1103–1113, 1989.

[54] P. Yu. Chebotarev, “Aggregation of preferences by the generalized row sum method”, *Mathematical Social Sciences*, Vol. 27, pp. 293–320, 1994.

[55] E. Shamis, “Graph-theoretic interpretation of the generalized row sum method”, *Mathematical Social Sciences*, Vol. 27, pp. 321–333, 1994.

[56] P. Yu. Chebotarev and E. Shamis, “On the proximity measure for graph vertices provided by the inverse Laplacian characteristic matrix”, in *Abstracts of the Fifth Conference of the International Linear Algebra Society*, (Atlanta), pp. 30–31, Georgia State University, 1995.

[57] P. Yu. Chebotarev and E. Shamis, “Characterizations of scoring methods for preference aggregation”, *Annals of Operations Research*, 1998. (Forthcoming).

[58] E. Zermelo, “Die Berechnung der Turnier-Ergebnisse als ein Maximumproblem der Wahrscheinlichkeitsrechnung”, *Math. Zeit.*, Vol. 29, pp. 436–460, 1928.

[59] R. A. Bradley and M. E. Terry, “The rank analysis of incomplete block designs. 1. The method of paired comparisons”, *Biometrika*, Vol. 39, pp. 324–345, 1952.

[60] M. Taylor, “Influence structures”, *Sociometry*, Vol. 32, pp. 490–502, 1969.

[61] N. E. Friedkin, “Theoretical foundations for centrality measures”, *American Journal on Sociology*, Vol. 96, pp. 1478–1504, 1991.